

**Collybia tuberosa****Braunknolliger Sklerotienröbling**

Fungi, Dikarya, Basidiomycota, Agaricomycotina, Agaricomycetes, Agaricomycetidae, Agaricales, Tricholomataceae

Legitimate (Bull.) P. Kumm. 1871

Aktueller Name gem. MycoDB: *Collybia tuberosa* (Bull.) P. Kumm., Der Führer in die Pilzkunde: 116 (1871) [MB242880]Basionym: *Agaricus tuberosus* Bull., Herbar de la France 6: t. 256 (1786) [MB154987]

Obligate Synonyme:

Microcollybia tuberosa (Bull.) Lennox, Mycotaxon 9 (1): 187 (1979) [MB300984]*Agaricus amanitae* subsp. *tuberosus* (Bull.) Pers., Observationes mycologicae 2: 53 (1800) [MB495119]

Taxonomische Synonyme:

Marasmius sclerotipes Bres., Fungi Tridentini 1 (1): 12, t. 11:1 (1881) [MB213222]*Collybia sclerotipes* (Bres.) S. Ito, Mycol. Fl. Japan: 123 (1950) [MB328717]*Chamaeceras sclerotipes* (Bres.) Kuntze, Revisio generum plantarum 3 (3): 457 (1898) [MB524386]*Gymnopus tuberosus* Gray, A natural arrangement of British plants 1: 611 (1821) [MB486482]Gelbknolliger Sklerotienröbling / *Collybia cookei*

makroskopisch

SubstratStark abgebauten *Lactarius* und *Russula***Gattung/en:**

Collybia

<https://www.mycopedia.ch/pilze/1015.htm>

Flammer, T©

16245 19.10.2023