



Exidia thuretiana

Weisslicher Drüsling

Fungi, Dikarya, Basidiomycota, Agaricomycotina, Agaricomycetes, Auriculariomycetidae, Auriculariales, Exidiaceae

Legitimate

(Lév.) Fr. 1874

Aktueller Name gem. MycoDB: Exidia thuretiana (Lév.) Fr., Hymenomycetes europaei: 694 (1874) [MB179018]

Basionym: Tremella thuretiana Lév., Annales des Sciences Naturelles Botanique 9: 127 (1848) [MB227814]

Taxonomische Synonyme: Heterochaete europaea Höhn., Annales Mycologici 1 (5): 393 (1903) [MB241334]

makroskopisch

Fruchtkörper / Habitus / Wachstumsform

Gelatinös, ohne Drüsenwärtchen, linsenförmig, dann zusammenfliessend, abgesetzte Wachstumsform

Substrat

Quercus

Fruchtkörperfarbe / Farbspektrum

Weisslich

mikroskopisch

Sporenform

Allantoid

Basidien

Hypobasidien septiert mit langen Epibasidien

Hyphen / Hyphensystem

Mit Schnallen

Gattung/en:

Exidia

<https://www.mycopedia.ch/pilze/3725.htm>



Exidia thuretiana

Weisslicher Drüsling

Fungi, Dikarya, Basidiomycota, Agaricomycotina, Agaricomycetes, Auriculariomycetidae, Auriculariales, Exidiaceae




Flammer, T©

12135 26.02.2021




Flammer, T©

12136 26.02.2021



Flammer, T©

12139 26.02.2021



Flammer, T©

12137 26.02.2021



Exidia thuretiana

Weisslicher Drüsling

Fungi, Dikarya, Basidiomycota, Agaricomycotina, Agaricomycetes, Auriculariomycetidae, Auriculariales, Exidiaceae

