

**Hemimycena candida****Beinwell-Scheinhelmling, Rasiger Scheinhelmling**

Fungi, Dikarya, Basidiomycota, Agaricomycotina, Agaricomycetes, Agaricomycetidae, Agaricales, Tricholomataceae

Legitimate

(Bres.) Singer 1943

Aktueller Name gem. MycoDB: *Hemimycena candida* (Bres.) Singer, *Annales Mycologici* 41: 121 (1943) [MB259584]Basionym: *Omphalia candida* Bres., *Fungi Tridentini* 2 (14): 87 (1900) [MB213980]

Obligate Synonyme:

Helotium candidum (Bres.) Redhead, *Canadian Journal of Botany* 60 (10): 2004 (1982) [MB109818]*Mycena candida* (Bres.) Kühner, *Encyclopédie Mycologique* 10: 659 (1938) [MB251231]*Delicatula candida* (Bres.) Kühner & Romagn., *Flore Analytique des Champignons Supérieurs*: 119 (1953) [MB296510]*Marasmiellus candidus* (Bres.) Singer, *Lilloa* 22: 299 (1951) [MB508844]*Omphalia lepista* var. *candida* (Bresadola) Quélet, *Enchiridion Fungorum in Europa media et praesertim in Gallia Vigentium*: 22 (1886) [MB803051]

makroskopisch

Fruchtkörper / Habitus / Wachstumsform

Büschelig bis rasig an den Wurzeln des Beinwells wachsend

Gattung/en:

Hemimycena

<https://www.mycopedia.ch/pilze/1036.htm>

Flammer, T©

10320 10.11.2019