



## Radulomyces molaris

Fungi, Dikarya, Basidiomycota, Agaricomycotina, Agaricomycetes, Agaricomycetidae, Agaricales, Radulomycetaceae, Radulomyces

Legitimate (Chaillet ex Fr.) M.P. 1960  
Christ.

Aktueller Name gem. MycoDB: Radulomyces molaris (Chaillet ex Fr.) M.P. Christ., Dansk Bot. Ark. 19 (2): 232 (1960) [MB338151]

Basionym: Radulum molare Chaillet ex Fr., Elenchus Fungorum 1: 151 (1828) [MB176952]

Obligate Synonyme:

Cerocorticium molare (Chaillet ex Fr.) Jülich & Stalpers, Verhandelingen Koninklijke Nederlandse Akademie van Wetenschappen Afdeling Natuurkunde 74: 72 (1980) [MB111209]

Basidioradulum molare (Chaillet ex Fr.) H. Furuk., Bulletin of the Government Forest Experimental Station Meguro 261: 59 (1974) [MB309458]

Taxonomische Synonyme:

Sistotrema rude Pers., Mycologia Europaea 2: 192 (1825) [MB440445]

Radulum rude (Pers.) S. Lundell, Fungi Exsiccati Suecici Fasc. 29-30: no. 1412 (1947) [MB282501]

Hydnum rude (Pers.) Duby, Botanicon gallicum 2: 780 (1830) [MB439966]

Sistotrema crustaceum Pers., Mycologia Europaea 2: 193 (1825) [MB440427]

mikroskopisch

### Sporenlänge

10 x 6,5 -7,5 µm

### Zystidenart

Leptocystiden: fusiform-subulat dicwandig, ca 35 µm lang

chemisch

### Melzers-Reagenz

IKI-

### Kalilaugenreaktion / KOH

Negativ

### Gattung/en:

Radulomyces

<https://www.mycopedia.ch/pilze/10307.htm>