



Sporormiella intermedia

Fungi, Dikarya, Ascomycota, Pezizomycotina, Dothideomycetes, Pleosporomycetidae, Pleosporales, Sporormiaceae

Legitimate (Auersw.) S.I. Ahmed & Cain ex Kobayasi 1969

Aktueller Name gem. MycoDB: Sporormiella intermedia (Auersw.) S.I. Ahmed & Cain ex Kobayasi, Bull. Tokyo Sci. Mus.: 339 (1969) [MB339558]

Basionym: Sporormia intermedia Auersw., Hedwigia 7: 67 (1868) [MB188603]

Obligates Synonym: Preussia intermedia (Auersw.) S. Ahmad, Monogr. Biol. Soc. Pakistan 8: 49 (1979) [MB320987]

makroskopisch

Fruchtkörper / Habitus / Wachstumsform

Perithezium schwarz mit verlängertem Hals, birnenförmig

Substrat

Kuhmist/Stroh

mikroskopisch

Sporenlänge

Reif 50 - 55 µm, unreif 41 - 45 µm x 8 - 10 µm

Sporenform

Unreif zylindrisch, reif eckig (die mittleren 2 Sporen), tief septiert, mit plus minus gerader Keimspalte

Sporenmerkmale

3fach Septiert

Ascus

Länge 200 µm

chemisch

Barals

Negativ

Varia

Bemerkungen / Hinweise / Abstract

Ellis: Ascospores deeply constricted at septa, easily fragmenting

Gattung/en:


Sporormiella

<https://www.mycopedia.ch/pilze/11494.htm>



Sporormiella intermedia


Fungi, Dikarya, Ascomycota, Pezizomycotina, Dothideomycetes, Pleosporomycetidae, Pleosporales, Sporormiaceae





Sporormiella intermedia

Fungi, Dikarya, Ascomycota, Pezizomycotina, Dothideomycetes, Pleosporomycetidae, Pleosporales, Sporormiaceae



Flammer, T©

14918 13.04.2023